

Supplemental Label



Arena® 50 WDG INSECTICIDE¹

EPA Reg. No. 59639-152
(Except California)

FOR ADDITIONAL USE DIRECTIONS ON RESIDENTIAL APPLE AND PEAR TREES (Not for use in commercial production sites)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Arena 50 WDG should be applied when threshold populations are observed. Do not wait until large insect populations have been established. Apply *Arena* 50 WDG in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for optimum control. To achieve optimum coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre.

RECOMMENDED FOLIAR POST BLOOM APPLICATION RATES FOR USE ON RESIDENTIAL BEARING APPLE AND PEAR TREES

Crop	Pests	<i>Arena</i> 50 WDG Application Rates	Special Instructions
Residential bearing: Apple Crabapple Pear	Aphids Leafhoppers Apple Maggot Leafminers Plum Curculio Codling Moth Leafrollers (suppression) Oriental Fruit Moth Soft Scales Pear Psylla	2 to 3 oz per 100 gallons of spray solution or 4 to 6 oz/A (2 to 3 oz ai/A) Rates per 100 gallon assume a spray rate of 200 gallons per acre. Adjust <i>Arena</i> 50 WDG rate per 100 gallons so as not to exceed 6 oz <i>Arena</i> 50 WDG per acre.	<i>Arena</i> 50 WDG should be applied when threshold populations are observed. Do not wait until large insect populations have been established. The amount of <i>Arena</i> 50 WDG required per acre will depend on tree size and pest pressure. Choose lower rates for light infestation and/or small trees, and the higher rates for heavy infestations and/or larger trees. Apply by ground in 200 gallons per acre as a full coverage spray. For optimum performance of Leafminers and leafrollers, make the first application as soon as pollination is complete and bees are removed from the orchard. Repeat application as necessary.

continued

RECOMMENDED FOLIAR POST BLOOM APPLICATION RATES FOR USE ON RESIDENTIAL BEARING APPLE AND PEAR TREES (continued)

- Do not apply treatments less than ten (10) days apart.
- Do not apply within seven (7) days of harvest.
- Do not apply more than 6.4 oz of *Arena* 50 WDG (0.2 lb ai) per acre per season.
- Do not feed or allow livestock to graze on cover crops from treated orchards.
- Applications must be made post bloom.

THIS SUPPLEMENTAL LABEL SUPERCEDES PREVIOUS LABELING WITH REVISED RATE STRUCTURE ON CERTAIN PESTS.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICATION DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Manufactured for
Valent U.S.A. Corporation
P. O. Box 8025
Walnut Creek, CA 94596-8025

Made in U.S.A.



Products That Work, From People Who Care® | www.valent.com | 800-89-VALENT (898-2536)

Read and follow the label instructions before using.

Products That Work, From People Who Care is a registered trademark of Valent U.S.A. Corporation. *Arena* is a registered trademark of Sumitomo Chemical Company, Limited. © 2008 Valent U.S.A. Corporation. All rights reserved. Printed in the USA.
2008-AREWDG-0012 EX CA 5/08 mf/AV